## **COVID-19 vaccination - what can you get?**



IF YOU ARE	YOU CAN GET
<ul> <li>Aged 75+ or you live in a care home for older adults</li> </ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>Booster from three months (91 days) after second dose</li> <li>Spring booster around 182 days (six months) after previous dose, unless invited earlier</li> </ul>
<ul><li>Aged 18 to 74</li></ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>Booster from three months (91 days) after second dose</li> </ul>
<ul><li>Aged 16 to 17</li></ul>	<ul> <li>Two doses, at least 12 weeks apart</li> <li>Booster from three months (91 days) after second dose</li> </ul>
<ul> <li>Aged 12+ and you are, or have been, immunosuppressed</li> </ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>If severely immunosuppressed at the time of earlier doses, a third primary dose, typically eight weeks after second dose (timing may vary and should be discussed with a clinician)</li> <li>Booster dose and spring booster from three months (91 days) after previous dose</li> </ul>
<ul> <li>Aged 12-15 and with a health condition that puts you at higher risk from COVID-19, or you are a household contact of someone who is immunosuppressed</li> </ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>Booster from three months (91 days) after second dose</li> </ul>
<ul><li>Aged 12-15 and not in any of the above groups</li></ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>Booster dose under review</li> </ul>
<ul> <li>Aged 5-11 with a health condition that puts you at higher risk from COVID-19, or you are a household contact of someone who is immunosuppressed</li> </ul>	<ul> <li>Two doses, at least eight weeks apart</li> <li>Booster dose under review</li> </ul>

To book your COVID-19 vaccination appointments visit www.nhs.uk/covid-vaccine or call 119. For information on walk-in sites available near you visit www.nhs.uk/vaccine-walk-in.